

Ecpyrrhorhoe diffusalis (Guenée, 1854) is a relict species in Hungary (Lepidoptera: Crambidae)

IMRE FAZEKAS

Regiograf Institute, Majális tér 17/A, H-7300 Komló, Hungary, e-mail: fazekas@micrilepidoptera.hu

FAZEKAS, I.: *Ecpyrrhorhoe diffusalis* (Guenée, 1854) is a relict species in Hungary (Lepidoptera: Crambidae).

Abstract: The postglacial relict species *Ecpyrrhorhoe diffusalis* (Guenée, 1854) is a very rare crambid species in Hungary. The distribution of this species in Hungary and Europe is reviewed based on collection and literature data.

Keywords: Lepidoptera, Crambidae, *Ecpyrrhorhoe diffusalis*, relict species, biology, distribution, Hungary.

Introduction

All available information on the distribution of *Ecpyrrhorhoe diffusalis* in Hungary and Europe is collected. The published data based on faunistic papers and lepidopterological collections. In the last years (1980-2012), the author intensively surveyed the Hungarian habitats of this moth. These data comprise 22 years field works.

Ecpyrrhorhoe diffusalis was described from France (GUENÉE 1854). So far, this species has been hardly studied. The distribution of this species is disjunct. The moth is relatively widespread but occurs always locally and rare from India and Asia Minor to southern Europe and Canary Island; from about 150 m up to 1600 m altitudes (SZENTIVÁNY & UHRIK 1942, KLIMESCH 1968, MATHEW 2006).

Taxonomy

Lepidoptera

Crambidae

Pyraustinae

Ecpyrrhorhoe Hübner, [1825], 1816

Harpadispar Agenjo, 1952

Pyraustegia Marion, 1963

Yezobotys Munroe & Mutuura, 1969,

Ecpyrrhorhoe diffusalis (Guenée, 1854)

Botys diffusalis Guenée, 1854, Historie naturelle de insects 8, p. 340. Locus typicus: "France méridionale, environs de Nîmes et Montpellier".

References: AGENJO 1952, BORHIDI 2003, DANIEL et al. 1951, FAZEKAS 1996, 2002, GUENÉE 1854, KLIMESCH 1968, MANN 1854, MAES 1994, MATHEW 2006, NUSS et al.

2012, OSTHELDER & PFEIFFER 1940, POPESCU-GORJ 1964, REBEL 1899, REBEL & ZERNY 1931, SLAMKA 2010, 2013, SZABÓKY 1980, 2000, SZENT-IVÁNY & UHRIK-MÉSZÁROS 1942.

Diagnosis: Variable species. Wingspan: 19-22mm. Forewings colouration brownish-ochreous or reddish; with more distinct ante- and postmedial lines. Antemedian line brownish white and wide, especially in middle. Postmedial line in middle part less bulging and continues on the hind wings where paler. Medial area darker with c-shaped distal spot. External area of wings slightly darker.

Distribution: According to NUSS et al. (2012), it's distribution in Europe is restricted to Albania, Bosnia and Herzegovina, Bulgaria, Canary Is, Croatia, France, Greek, Hungary, Italia, Macedonia, Romania, Spain, and Switzerland. Known outside of Europe: Morocco, Turkey, Syria, Transcaucasia, Iran, Afghanistan, Pakistan, the United Arab Emirates and India (SLAMKA 2013).

One old record for Western Ukraine (Lvov) is not confirmed by recent data, probably it is erroneous or misidentified (see SLAMKA 2010). *E. diffusalis* is apparently a very rare and local species in central and Mediterranean Europe. It is not present on Mediterranean islands.

Biology: No detailed information is available on habitat preference. Adults attracted to light or flying in the daytime between plants in Hungary. SZABÓKY (2000) wrote “*Differentially from the related Crambidae species, it flies only in daytime. So far, it has been collected by light only*”. This is a misstatement. In the last years (2000-2012), I intensively surveyed the moth's habitats and found thought the light-trap and light attracts the moth but it flies daytime as well. It is probably bivoltine, flies from May to mid-September. According to the literature the larva oligophagous on *Lavatera* and *Marrubium* species.


Investigation in Hungary

SZENT-IVÁNY and UHRIK-MÉSZÁROS (1942) published it from several localities outside the present Hungary: „Herkulesfürdő“ (in Romania: Baile Herculeană) and Zengg (in Croatia: Senj). SZABÓKY (1980) recorded firstly from Hungary: South Hungary, Villány Hills, Nagyharsány, near Croatian border.

The localities of *E. diffusalis* are situated in 190 m altitude and the habitats are typically calcareous open rock grasslands with numerous endemic and relict plant species (e.g. *Trigonella gladiata*, *Colchicum hungaricum*, *Medicago orbicularis*, *Orobanche nana*, *Sempervivum tectorum*). The characteristic association is *Sedo sopianae-Festucetum dalmaticae* on Triassic and Jurassic limestone SIMON 1964. The following other rare micro-moth species were also captured here: *Hepialus amasinus*, *Ecpyrrhorhoe diffusalis* and *Jordanita fazekasi*.

The isolated, relict and xerothermophilous *Ecpyrrhorhoe diffusalis* population in southern Hungary lives dominantly in this protected area (Natura 2000 site).

Collected data from Hungary: 2 males, “Villányi hg., Szársomlyó, 1979.VI.24. leg. Szabóky Cs., in coll. HNHM Budapest and Szabóky Cs.; 3 ex, Szársomlyó, szoborpark, 2003.VII.22. leg. Szabóky Cs. et Takács A., in coll. Buschmann F., Jászberény.



Figs. 1-4: Adult and habitat of *Ecpyrrhorhoe diffusalis* in Hungary:
1) adult, Villány Hills, Szársomlyó hill; 2) habitat in Szársomlyó hill;
3) adult, Dombóvár-Gunaras; 4) habitat in Dombóvár-Gunaras.




Fig. 5: The distribution of relict *Ecpyrrhorhoe diffusalis* in Hungary. Distribution of calcareous open rocky grasslands in country (grey colour): the potential range of species.

New results: 1-1 males, H-Nagyharsány, Szársomlyó, 230 m (N45°51'22,38"; E18°25'3,91"), 2000.07.02; 2003.07.12; 2010.08.; 2012.07.18., leg. Fazekas I., in coll. Regiograf Institute, H-Komló; Finally, Dombóvár: Gunaras is a very surprising new occurrence in south Hungary (outside the protected area). Dombóvár, Gunaras, 1 male, 20.07.2007, leg. et in coll. A. Schreurs (NL-Kerkrade), det. J. Asselbergs, revid. I. Fazekas. The species is new for Tolna county and was unknown there to 2012.




Fig. 6: Dot map of the distribution data of *Ecpyrrhorhoe diffusalis* southeast in Europe was made on the basis of collection and literature data (schematic).

Discussion

Ecpyrrhorhoe diffusalis is very rare but the Hungarian Red Data Book does not mention this (RAKONCZAY 1989), and does not protected by law. According to author, *Ecpyrrhorhoe diffusalis* is a regressive postglacial relict element.

Many micro-moths exist only in isolated colonies as relict populations, separated from their relatives by climatic or ecological effects in the Pannonian region. Their unique relict populations located in Hungarian middle hills and in the Great Hungarian Plain are typically placed on limestone or dolomite mountains and sand-hill areas.

The Villány Hills, with xerothermophilous post-glacial vegetation, certainly provides optimal conditions for the Mediterranean and for other sub-Mediterranean fauna elements. The populations of *Ecpyrrhorhoe diffusalis* restricted to limited areas in Hungary would deserve greater attention of the nature conservation authorities. I am working on a proposal to protect the habitats of this moth.

Acknowledgements

We thank to A. Schreurs (NL-Kerkrade) for providing data of this species from Dombóvár (Tolna county).

References

- AGENJO, R. 1952. Fáunula Lepidopterológica Almeriense. – Madrid: pp.1-370, pis. 1-24.
- BORHIDI, A. 2003: Magyarország növénytársulásai. – Akadémiai Kiadó, 610 p.
- DANIEL, F., FORSTER, W. & OSTHEIDER, L. 1951: Beiträge zur Lepeidopterenfauna Mezedoniens. – Veröffentlichungen der Zoologischen Staatssammlung München 2: 1-78.
- FAZEKAS, I. 1996: Systematic catalogue of the Pyraloidea, Pterophoridae and Zygaenoidea of Hungary (Lepidoptera). – Folia Comloensis, Suppl.: 1-34.
- FAZEKAS, I. 2002: Baranya megye Microlepidoptera faunájának katalógusa (Lepidoptera). – Folia Comloensis 11: 5-76.
- GUENÉE, A. 1854: Historie naturelle de insects 8. – Deltoides et Pyralites, Paris, pp. 1-448.
- KLIMESCH, J. 1968: Die Lepidopteren fauna Mezedoniens IV. Microlepidoptera. – Prirodonaucen Muzej Skopje, Posebno Izdanje No. 5, 202 p.
- MANN, J. 1854: Aufzählungen der Schmetterlinge, gesammelt auf einer Reise im Auftrage des k. k. Zoologischen Museums nach Oberkrain und dem Küstenlande, in den Monaten Mai und Juni 1854, als Beitrag zur Fauna des österreichischen Kaiserstaates. – Verhandlungen der Zoologisch-Botanischen Gesellschaft in Wien 4: 515-596.
- MAES, K. V. N. 1994. Some notes on the taxonomic status of the Pyraustinae (sensu Minet 1981 [1982]) and a check list of the Palearctic Pyraustinae (Lepidoptera, Pyraloidea, Crambidae). – Bulletin et Annales de la Societe Royale Belge d'Entomologie 130: 159-168.
- MATHEW, G. 2006: An inventory of Indian Pyralids (Lepidoptera: Pyralidae). – Zoos' Print Journal 21 (5): 2245-2258.
- NUSS, M., SPEIDEL, W. & SEGERER, A. 2012: Ecpyrrhorhoe diffusalis (Guenée, 1854). – Fauna Europaea. <http://www.faunaeuropaea.org> (Accessed: 22.08.2012)
- OSTHEIDER, L. & PFEIFFER, E. 1940: Lepidopteren-Fauna von Marasch in türkisch Nordsyrien. – Mitteilungen der Münchener Entomologisches Gesellschaft 30: 107-116.
- POPESCU-GORJ, A. 1964: Catalogue de la collection de Lépidoptères „Prof. A. Ostrogovich” du Muséum D'histoire Naturelle „Grigore Antipa” Bucarest. – Bucarest, 293 p.
- RAKONCZAY, Z. (ed.) 1989: Vörös könyv [Hungarian Red Data Book]. – Akadémiai Kiadó, Budapest, 360 p.
- REBEL, H. 1899: Zweiter Beitrag zur Lepidopteren-Fauna Südtirols. – Verhandlungen der Kaiserlich-Königlichen Zoologisch-Botanischen Gesellschaft in Wien, 1899: 158-185.
- REBEL, H. & ZERNY, H. 1931: Die Lepidopterenfauna Albaniens. – Denkschriften Akademie der Wissenschaften in Wien, Mathematische-naturwissenschaftliche Klasse 103: 38-161.
- SLAMKA, F. 2010: Pyraloidea (Lepidoptera) of Central Europe. – František Slamka, Bratislava, 176 p.
- SLAMKA, F. 2013: Pyraloidea of Europe (Lepidoptera) Volume 3, Pyraustinae & Spilomelinae. – František Slamka, Bratislava, 357 p.
- SZABÓKY, Cs. 1980: Magyar faunára új molylepkek (Lepidoptera). – Folia Entomologica Hungarica 33: 204-208.
- SZABÓKY, Cs. 2000: A Villányi-hegység molylepkei (Microlepidoptera). – Dunántúli Dolgozatok Természettudományi Sorozat, Pécs, 10: 297-307.
- SZENT-IVÁNY, J. & UHRÍK-MÉSZÁROS, T. 1942: Die verbreitung der Pyralididen (Lepidopt.) im Karpatenbeckens. – Annales Historicoo-naturalis Musei Nationalis Hungarici, Pars Zoologica, 35: 105-196.